



Ocular and Systemic Complications of Type II Diabetes: Sociodemographic Predictors of Emergency Department Utilization



Chaz Crosby¹, Dr. Ronald Horswell², San Chu², Grace Kim³, Dr. Lucio Miele³

¹Tulane University School of Medicine, New Orleans, LA

²Pennington Biomedical Research Center, Baton Rouge, LA

³Louisiana State University Health Sciences Center, New Orleans, LA

Introduction

INTRODUCTION:

- Lack of access to primary care and lack of continuity of care-----> Increased use of emergency department (ED) when experiencing chronic disease symptoms
- Structural barriers (systemic racism, insurance instability, socioeconomic challenges) further exacerbate ED use and disease management
- Poorly managed type II diabetes ----> Development of ocular and systemic complications

BACKGROUND:

- Diagnosed Type II Diabetes Statistics in the United States***
 - 38 million Americans have diabetes, with 90-95% having type II diabetes (CDC, 2024)
 - However, many more Americans have diabetes and are undiagnosed
 - Diabetic complications include ocular (retinopathy, cataracts), kidney disease, nerve damage, foot ulcers, heart disease, amputation risks and more (CDC, 2024)
- Healthcare Costs & the ED***
 - 18 million preventable ED visits adds 32 billion dollars to the healthcare system each year (UnitedHealth Group, 2019)
- Differential Outcomes of Type II Diabetes***
 - Socioeconomic status, neighborhood & built environment, food access, healthcare access contribute to varied outcomes for patients with diabetes (Hills-Briggs et al., 2020)
- Gap in Literature***
 - Few studies jointly evaluate diabetes complications and ED visits by sociodemographic factors

PROJECT AIM:

- Examine how the ED is utilized to manage type II diabetes complications, evaluated through a sociodemographic lens


OBJECTIVES AND METHODOLOGY:

- Data collection: All of Us Workbench- NIH funded program that prioritizes research from 900,000 participants from diverse and historically underrepresented populations
 - R computational tool
- Establish the rate of ED utilization by diabetic complication stratified by self-reported race
- Analyze the representation of self-reported race and insurance status in diabetic complications
 - Insurance status will serve to estimate the role of SES in racial disparities

Average ED Visits for Type II Diabetic Complications Stratified by Self-Reported Race

Figure 1		
Complication	Black Patients	White Patients
Retinopathy / Cataracts	≈21 avg ED visits SE = 0.84 n = 1670	≈14 avg ED visits SE = 0.59 n = 2119
Foot Ulcers	≈29 avg ED visits SE = 1.99 n = 534	≈17 avg ED visits SE = 1.11 n = 1077
Kidney Dysfunction	≈23 avg ED visits SE = 1.05 n = 2871	≈14 avg ED visits SE = 0.44 n = 4464

Patients with "Diabetes with Complications" Stratified by Self-Reported Race and Insurance Status



Results

Figure 1:

- In each complication examined (retinopathy/cataracts, foot ulcers, and kidney dysfunction), Black patients experienced more ED visits than White Patients, reflecting systemic inequities in primary and preventative care

Figure 2:

- Black patients were disproportionately represented in both Medicaid (44%) and private insurance (26%) groups for the "Diabetes with Complications" group, relative to the other racial groups

Conclusion

- ED utilization varies based on diabetic complication, sociodemographic factors, and access to care
- Black patients were overrepresented in Medicaid and private insurance groups, suggesting systemic barriers beyond insurance type
 - Are some groups at greater risk of developing complications, driving higher ED utilization?*
 - Something else to consider:* Patients on Medicaid oftentimes have limited access to primary care (reluctance to accept Medicaid coverage), reduced financial disincentives, and higher chronic disease burden. *Could this be an additional factor in increased ED use?*
- Findings will help identify at-risk populations and inform targeted interventions to reduce ED visits, optimize diabetes management, and improve health system efficiency
- Next steps:**
 - Additional ED visit analyses
 - Risk Score Development
 - Patient Engagement Program

References

Centers for Disease Control and Prevention. (2024, May 15). *Type 2 diabetes*. National Center for Chronic Disease Prevention and Health Promotion. <https://www.cdc.gov/diabetes/about/about-type-2-diabetes.html>

Centers for Disease Control and Prevention. (2024, May 15). *Diabetes complications*. National Center for Chronic Disease Prevention and Health Promotion. <https://www.cdc.gov/diabetes/complications/index.html>

UnitedHealth Group. (2019, July). *Avoidable emergency department visits: An analysis of factors and strategies* [PDF]. UnitedHealth Group. <https://www.unitedhealthgroup.com/content/dam/UHG/PDF/2019/UHG-Avoidable-ED-Visits.pdf>

Hill-Briggs, F., Adler, N. E., Berkowitz, S. A., Chin, M. H., Gary-Webb, T. L., Navas-Acien, A., Thornton, P. L., & Haire-Joshu, D. (2020). Social Determinants of Health and Diabetes: A Scientific Review. *Diabetes care*, 44(1), 258–279. Advance online publication. <https://doi.org/10.2337/dci20-0053>