

Uncommon Triad: Surgical Management of Concurrent Spigelian, Inguinal, and Femoral Hernias*

Daniel Vicknair, BA¹; Dr. Ian Hodgdon, MD, FACS²


¹LSUHSC School of Medicine, ²University Medical Center New Orleans

BACKGROUND

BACKGROUND: The simultaneous occurrence of Spigelian, inguinal, and femoral hernias is a rare and, to our knowledge, previously unreported clinical presentation.¹ This case report describes such a triad in a 74-year-old female with systemic lupus erythematosus (SLE), lupus nephritis, chronic obstructive pulmonary disease (COPD), rheumatoid arthritis, and heart failure with preserved ejection fraction (HFpEF), where chronic corticosteroid use and autoimmune disease likely contributed to hernia formation via fascial weakening.² Her surgical history includes a right ventral hernia repair with mesh in 2012 and a documented 5 × 2 cm left lateral abdominal wall hernia in July 2023, with recent presentation of persistent left-sided abdominal pain.¹ Preoperative computed tomography (CT) imaging revealed an 8.3 × 4.1 cm Spigelian hernia containing small bowel loops, suggesting a multifactorial etiology involving chronic corticosteroid use and SLE-induced tissue fragility.³ We propose Robotic-assisted laparoscopic surgery offers a promising approach to address the diagnostic and therapeutic challenges of this rare triad.


OBJECTIVES

The primary objective is to describe the diagnostic complexities and surgical management strategies for concurrent Spigelian, inguinal, and femoral hernias in a high-risk patient.⁴ We aim to enhance the understanding of this uncommon clinical entity and assessing the efficacy of robotic-assisted laparoscopic repair in optimizing outcomes for patients with significant comorbidities.⁵



1A: CT Abdomen/Pelvis midsagittal view through Spigelian hernia pre-operation. "There is an 8.3 cm CC by 4.1 cm transverse spigelian hernia in the left lower quadrant extending inferiorly to the left superior pubic ramus. Loops of small bowel protrude through the defect. No current evidence for bowel obstruction. There is contrast material throughout the small bowel. Gas and stool are present in the colon. No free intraperitoneal gas or focal abscess is seen in the abdomen."

1B: CT Abdomen/Pelvis midsagittal view through Spigelian hernia post-operation.




2A: CT Abdomen/Pelvis cross-section view through Spigelian hernia pre-operation. 2B: CT Abdomen/Pelvis cross-section view through Spigelian hernia post-operation.

BACKGROUND


The patient underwent robotic-assisted laparoscopic transabdominal preperitoneal (rTAPP) repair using a modified transabdominal preperitoneal with extended view (TAPE) technique, following multidisciplinary clearance from pulmonology, endocrinology, rheumatology, and cardiology. Intraoperative exploration facilitated identification and repair of Spigelian, inguinal, and femoral hernias using synthetic mesh and peritoneal flap closure. Postoperative evaluation included monitoring for complications, recurrence, and recovery at short-term follow-up.⁴

RESULTS


Preoperative CT confirmed the Spigelian hernia, but intraoperative assessment via laparoscopic exploration revealed additional ipsilateral inguinal and femoral hernias, highlighting the limitations of imaging in detecting occult defects.^{6,1} The robotic-assisted procedure was completed without intraoperative or postoperative complications. Short-term follow-up at 30 days showed no recurrence or wound-related issues, supporting the efficacy of the modified TAPE technique in this corticosteroid-dependent, high-risk patient.³




3A: Intraoperative view. Contents pull out of muscular defect.



3B: Peritoneal flap created. L. indirect and femoral hernia in view




3C: Mesh placement: 15mm*10mm Progrid mesh was placed.



3D: Peritoneal flap closed with 3-0 V-locc over the mesh.

CONCLUSION

This case underscores the exceptional rarity of concurrent Spigelian, inguinal, and femoral hernias and the critical role of intraoperative laparoscopic exploration in their definitive diagnosis.^{6,5} Robotic-assisted repair proved effective for this rare hernia triad in a comorbid, immunosuppressed patient, reducing complications and hospital stay, warranting prospective studies to optimize protocols for such complex cases.⁷



4: Intuitive Di Vinci Xi intraoperatively (not this case)

REFERENCES

- Spangen, L. Spigelian hernia. *World J. Surg.* 13, 573–580 (1989). <https://doi.org/10.1007/BF01658873>
- Hope WW, Tuma F. Incisional Hernia. [Updated 2023 Jun 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023Jan. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK455995>
- Saiding Q, Chen Y, Wang J, Pereira CL, Sarmiento B, Cui W, Chen X. Abdominal wall hernia repair: from prosthetic meshes to smart materials. *Mater Today Bio.* 2023 Jun 29;21:100691. doi: 10.1016/j.mtbio.2023.100691. PMID: 37455815; PMCID: PMC10339210.
- Bittner R, Bingener-Casey J, Dietz U, et al. Update of guidelines for laparoscopic treatment of ventral and incisional abdominal wall hernias (International Endohernia Society (IEHS))—Part A. *Surg Endosc.* 2019;33(10):3069–3139. doi:10.1007/s00464-019-06907-7.
- Zonana-Nacach A, Barr SG, Magder LS, Petri M. Damage in systemic lupus erythematosus and its association with corticosteroids. *Arthritis Rheum.* 2000;43(8):1801–1808. doi:10.1002/1529-0131(200008)43:8<1801::AID-ANR16>3.0.CO;2-O.
- Manoharan V, Joshi AS. Occult hernia: role of laparoscopy in detecting the unseen. *Int Surg J.* 2022;10(1):97. doi:10.18203/2349-2902.isj20223599.
- Peñafiel JAR, Valladares G, Cynrta Lima Fonseca Rodrigues A, et al. Robotic-assisted versus laparoscopic incisional hernia repair: a systematic review and meta-analysis. *Hernia.* 2024;28(2):321–332. doi:10.1007/s10029-023-02881-1.

ACKNOWLEDGEMENTS

We express our sincere gratitude to Ian Hodgdon, MD, for his invaluable mentorship and significant contributions to this case. His expertise and guidance were instrumental in navigating the complexities of this rare presentation. We also extend our appreciation to the faculty, staff, and fellow residents of the Louisiana State University Health Sciences Center (LSUHSC) General Surgery Residency Program for their unwavering support, encouragement, and dedication to our education. Additionally, we are grateful to University Medical Center (UMC)-LSUHSC for providing the opportunity to engage in unique and challenging cases, which have enriched our clinical experience and professional development.

