

# Ankle Closure Methods: An Expert Survey of Orthopedic Trauma Association (OTA) Members and Review of the Literature



Jack McKay MD<sup>1</sup>, Christopher Bloise<sup>2</sup>, Claudia Leonardi PhD<sup>3</sup>, Peter Krause MD<sup>1</sup>

<sup>1</sup>Department of Orthopaedics, Louisiana State University Health Sciences Center School of Medicine, New Orleans, LA, USA

<sup>2</sup>Louisiana State University Health Sciences Center School of Medicine, New Orleans, LA, USA

<sup>3</sup>Behavioral and Community Health Sciences, School of Public Health, Louisiana State University Health Sciences Center, New Orleans, LA, USA

## Introduction

Ankle fractures are very common, accounting for 10.2% of all bone injuries, with an incidence in adults of 179 fractures per 100,000 persons per year (1,2). Ankle fractures are mostly caused by falls (61%) followed by sports injuries (22%) and other trauma mechanisms (17%) (1). Depending on the type of injury, ankle fractures can be treated either operatively or nonoperatively. Due to it being associated with superior outcomes though, surgery, particularly open reduction and internal fixation (ORIF), is considered the predominant treatment method for ankle fractures (4).

Wound complication rates after ORIF of ankle fractures vary between 1.4% and 18.8% (5). One of the most prevalent complications, surgical site infections (SSI), have been shown to negatively affect bony union and functional recovery of the ankle (4). Patient factors, such as a history of diabetes, peripheral neuropathy, and medications, are significantly associated with ankle fracture surgery wound complications (5). Although there is substantial evidence indicating skin closure methods are significantly associated with wound complications and surgical outcomes in various types of surgeries, to the best of our knowledge, only one study with a small sample size has investigated this association following ankle fracture surgery, yielding insignificant results (6).

Because of the lack of the subcutaneous support around the ankle, skin closure following ankle surgery is particularly challenging (6). Due to this unique challenge and the scarcity of literature examining skin closure methods following ankle surgery, we sought to evaluate the preferences of orthopedic trauma experts regarding skin closure methods after ORIF of rotational ankle fractures and compare these preferences to the current literature on outcomes of common skin closure methods.

## Materials and Methods

A 23 item web-based questionnaire was advertised to active members of the OTA from January to September of 2017. The survey was completed by 167 respondents. 150 (92%) of respondents were male, and the mean age was 45.1 years. The mean number of years in practice was 12.19 and ranged from 0-44 years (Table 1). The questionnaire was designed by the authors. The survey responses were recorded using the LSUHSC Redcap database. Using a cross-sectional survey study design, we evaluated the preferences of suture type and technique for skin closure after fixation of rotational ankle injuries. The majority of respondents are fellowship trained (147, 89.6%), and practice in a Level 1 academic setting (99, 60%). 96 (57.8%) of respondents have an academic practice. 48 (28.9%) reported being in private practice, while the remaining (22, 13.3%) chose "other" as their practice type.


A literature review was also conducted using the main medical databases to evaluate the available literature and to analyze the results of studies examining outcomes following common skin closure methods in a variety of surgeries. A total of 41 studies examining wound closure methods were identified. 17 were excluded due to reasons that included very small samples sizes, too specific study populations that results would not be able to be applied to our study, and studies that examined adhesive closure, leaving 24 studies that were included in our review. Five compared suture types and twenty compared sutures versus staples. Nine were randomized controlled trials, five were retrospective cohorts, and ten were systematic reviews/ meta-analyses. Nine studies specifically looked at orthopaedic surgeries.

## Results


167 of 1639 OTA members with either "Active," "Clinical," "Associate," or "Candidate" status responded to the survey for a response rate of 10.2%. Nearly all respondents indicated that they close these wounds in a layered fashion (90.4%). The most utilized skin closure method after ORIF of routine ankle injuries was interrupted non-absorbable nylon suture (49.7%) (Figure 1). This preference for interrupted non-absorbable suture increases with treating patients thought to have a higher risk of wound complications (eg diabetics). 68.7% of respondents used interrupted non-absorbable sutures when closing skin for those they considered "high risk" (ie diabetics) (Figure 2). When used, non-absorbable sutures are most commonly removed in the 2 to 3 week post operative period (125, 77.6%) (Figure 3). When asked "Approximately what percentage operative ankle injuries in your practice experience some surgical incision complication?", 82 (49.7%) responded that these occurred in 1-5% of patients (Figure 4). In regards to how much time, in minutes, was spent closing bimalleolar ankle fractures 80 (48.2%) estimated 5-10 minutes, with 80 (48.2%) estimating >10 minutes (Figure 5). As seen from response to the survey by OTA members, there are multiple common and accepted methods of wound closure after ORIF of rotational ankle fractures.

Of the studies examining sutures vs staples, seven favored sutures and two favored staples when looking at wound complications. Of these studies, three meta-analysis examined orthopaedic surgeries with only one revealing sutures has less complications than staples. While there is conflicting evidence on which method produces less complications, staple closure was significantly associated with greater reported pain and worse cosmetic scores compared to sutures, but had the benefit of quicker closing time. When comparing suture methods, interrupted sutures were associated with a significantly less incidence of wound dehiscence compared to continuous sutures in two separate meta-analyses. Absorbable sutures were associated with significantly better cosmetic outcomes and no significant difference in overall wound complication rates compared to nonabsorbable sutures.


**Figure 1:** What is your primary method of skin closure for a routine ankle fractures? (N = 165).




**Figure 3:** If removable sutures are used, at what time point post operatively do you typically remove sutures if there are no wound complications? (N = 161).




**Figure 2:** What is your primary method of skin closure for a routine ankle fractures in high-risk patients (e.g. diabetics)? (N = 166).




**Figure 4:** Approximately what percentage operative ankle injuries in your practice experience some surgical incision complication? (N = 165).



**Figure 5:** Approximately how long, in minutes, do you spend closing surgical incisions for fixation of bimalleolar ankle fractures? (N = 166).



**Figure 6:** Who performs the majority of your closure? (N = 166).



## Conclusions

The best method of skin closure after ORIF of rotational ankle injuries remains unproven. The relative majority of OTA members prefer to close skin with interrupted nylon after a layered closure. The popularity of interrupted non-absorbable suture increased by 20% when comparing "routine ankle fractures" and "routine ankle fractures in high risk patients." This would suggest that most surgeons rely on this method of closure as the "safest" method. This would imply that most surgeons believe the interrupted non-absorbable closure method is most likely to prevent post operative wound complications. This method has also been shown to be associated with greater cosmetic scores and less pain compared to staples. The decreased rates of dehiscence compared to continuous/ running sutures seen in other surgery types support the higher usage rate following ORIF of ankle fractures.

## References

1. Elisei, R., Ogasawara, S. E., & Larson, P. (2018). Population-based epidemiology of 9787 ankle fractures. *Foot and Ankle Surgery*, 24(1), 34-39.
2. Jato, N., Nilsson, H., & Manktelow, P. (2018). Epidemiology of Ankle Fractures: 1765 cases identified in Northeastern County during 2009-2013 and classified according to AO/OTA. *BMC musculoskeletal disorders*, 19(1), 441.
3. Ramelli, S. (2016). Management of ankle fractures in the elderly. *EDOT open reviews*, 1(1), 239-246.
4. Sun, Y., Wang, H., Tang, Y., Zhao, H., Qin, S., Xu, L., & Zhang, F. (2018). Incidence and risk factors for surgical site infection after open reduction and internal fixation of ankle fracture: A retrospective multicenter study. *Medicine*, 97(17).
5. Miller, A. G., Margolis, A., & Raskin, S. M. (2012). Risk factors for wound complications after ankle fracture surgery. *JBS*, 94(22), 2047-2052.
6. Lehtonen, E., Patei, H., Phillips, S., Pinto, M. C., Narasim, S., & Shah, A. (2018). Staple versus suture closure for ankle fracture fixation: Retrospective chart review for safety and outcomes. *The Foot*, 37, 71-76.
7. Tsujinaka, Y., Yamamoto, K., Fujita, J., Endo, S., Kawada, J., Nakahira, S., ... & Mori, M. (2013). Subcuticular sutures versus staples for skin closure after open gastrointestinal surgery: a phase 3, multicentre, open-label, randomised controlled trial. *The Lancet*, 382(9989), 1305-1312.
8. Imamura, K., Adachi, K., Sasaki, R., Momma, S., Shinoh, S., Seyama, Y., ... & Kaneko, T. (2016). Randomized comparison of subcuticular sutures versus staples for skin closure after open abdominal surgery: a multicenter open-label randomized controlled trial. *Journal of Gastrointestinal Surgery*, 20(12), 2083-2092.
9. Kobayashi, S., Ito, M., Yamamoto, S., Kinugasa, Y., Kotaka, M., Saida, Y., ... & Moriya, Y. (2015). Randomized clinical trial of skin closure by subcuticular suture or skin stapling after elective colorectal cancer surgery. *Journal of British Surgery*, 20(5), 495-500.
10. Figueroa, D., Jank, V. C., Strydomski, A., Garner, R., Biggio, J. R., Andrews, W. W., ... & Tita, A. T. (2013). Surgical staples compared with subcuticular suture for skin closure after cesarean delivery: a randomized controlled trial. *Obstetrics and gynecology*, 121(1).
11. Bai, M., Zheng, X., Sun, S., Li, C. Y., Zhang, M. C., Guo, K. J., ... & Pang, Y. (2018). A prospective randomized comparison of 2 skin closure techniques in primary total hip arthroplasty surgery. *Hip International*, 28(1), 101-105.
12. Kuraki, I., M., Mullen, M., Masada, L. S., Wu, H., Liu, J., Mutch, D. G., ... & Novetsky, A. P. (2017). Wound complication rates after staples or suture for midline vertical skin closure in obese women: a randomized controlled trial. *Obstetrics and gynecology*, 129(1), 91.
13. Mackeen, A. D., Khalifeh, A., Fehse, J., Vogell, A., Han, C., Sendecki, J., ... & Bergheva, V. (2014). Suture compared with staple skin closure after cesarean delivery: a randomized controlled trial. *Obstetrics & Gynecology*, 123(6), 1169-1175.
14. Yuenyongwat, V., K., Iamthongpon, T., Hongnapanak, B., Tangtrakulwanich, "A Randomised Controlled Trial Comparing Skin Closure in Total Knee Arthroplasty in the Same Knee: Nylon Sutures versus Skin Staples." *Bone and Joint Research* 5 (2016): 185-190. Published Web 28 Oct. 2016.
15. Oweid, S., Borik, R., Rhoads, H., Jadhav, A., Soriano, S., & Oweid, R. (2017). Surgical staples: a superior alternative to sutures for skin closure after neck dissection—a single-blinded prospective randomized clinical study. *Journal of Oral and Maxillofacial Surgery*, 75(12), 2707-e1.
16. Hrabek, H., Wastler, F., Hahn, J., & Salomon, D. (2014). Metal staples versus conventional suture for wound closure in total knee arthroplasty. *Acta chirurgica orthopaedica et traumatologica czechoslovaca*, 81(3), 233-237.
17. Eggers, M. D., Fang, L., & Liorberger, D. R. (2011). A comparison of wound closure techniques for total knee arthroplasty. *The Journal of orthopaedics*, 26(8), 1253-1258.
18. Schröder-Pedanz, T. L., Ruck, M. P., Kalkas, S., Tomasiak, C., Nigro, S. D., & Yeiser, J. D. (2016). Incidence of wound complications in cesarean deliveries following closure with absorbable subcuticular staples versus conventional skin closure techniques. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 206, 53-56.
19. Chhabra, S., Gonnella, M., Sugimoto, M., Nomura, S., Kobayashi, S., Takahashi, S., ... & Konishi, M. (2018). Absorbable skin closure using subcuticular sutures prevents incisional surgical site infection in hepatopancreatobiliary surgery. *Surgery*, 164(2), 251-256.
20. Tuuli, M. G., Renspers, R. M., Carbone, J. F., Smailio, D., Macones, G. A., & Dalbo, A. D. (2011). Staples compared with subcuticular suture for skin closure after cesarean delivery: a systematic review and meta-analysis. *Obstetrics & Gynecology*, 117(3), 682-689.
21. Wang, H., Hong, S., Teng, H., Qiao, L., & Yin, H. (2016). Subcuticular sutures versus staples for skin closure after cesarean delivery: a meta-analysis. *The Journal of Maternal-Fetal & Neonatal Medicine*, 29(22), 3705-3711.
22. Lu, Y., Wang, C., Lin, L., Qin, Q., & Li, Q. (2018). Complication rate of different wound closures after primary hip arthroplasty-A survey of 373 patients. *Asian-Pacific journal of sports medicine, orthopaedics, rehabilitation and technology*, 11, 15-18.
23. Krolnick, R., Mackinnon, S. G., & Makandiah-Mehra, M. S. (2016). Comparing sutures versus staples for skin closure after orthopaedic surgery: systematic review and meta-analysis. *BMC open*, 6(1), e009257.
24. Kim, K. Y., Anouchiravan, A. A., Long, W. J., Vygoloch, J. M., Fernandez-Madrid, I., & Schwarzkopf, R. (2017). A Meta-Analysis and Systematic Review Evaluating Skin Closure After Total Knee Arthroplasty—What is the Best Method? *The Journal of orthopaedics*, 22(8), 2920-2927.
25. Hemming, K., Pinkney, T., Futaba, K., Pennington, M., Morton, D. G., & Lifford, R. J. (2013). A systematic review of systematic reviews and panoramic meta-analysis: staples versus sutures for surgical procedures. *PLoS one*, 8(10), e75332.
26. Smith, T. O., Senn, D., Mann, C., & Donell, S. (2010). Sutures versus staples for skin closure in orthopaedic surgery: meta-analysis. *Bmj*, 340, c1399.
27. Iarazzo, C., Gekles, I. D., Vouyoumou, E. K., Mamalis, I., Peppas, G., & Falagas, M. E. (2011). Sutures versus staples for the management of surgical wounds: a meta-analysis of randomized controlled trials. *The American Surgeon*, 77(9), 1206-1212.
28. McLean, R. A., H. B. Frye, F. F. First, B. H. Irvine, and M. H. Calvert. "Comparison of Skin Closure Using Continuous and Interrupted Nylon Sutures." *British Journal of Surgery* Br. J. Surg. 67 9 (1980): 633-35. Web.
29. Gurutunay, K. S., Toon, C. D., Allen, V. B., & Davidson, B. R. (2014). Continuous versus interrupted skin sutures for non-obstetric surgery. *Cochrane Database of Systematic Reviews*, (2).
30. Gupta, H., Vinayak, A., Menon, G. R., Agrawal, C. S., Chumber, S., & Kumar, S. (2008). Comparison of interrupted versus continuous closure in abdominal wound repair: a meta-analysis of 23 trials. *Asian journal of surgery*, 31(1), 104-114.
31. Xu, B., Yu, B., Wang, L., Chen, C., Yimaz, T. U., Zheng, W., & He, B. (2016). Absorbable versus nonabsorbable sutures for skin closure: a meta-analysis of randomized controlled trials. *Annals of plastic surgery*, 76(5), 598-606.