


Macy Owens<sup>1</sup>, Dean Smith, PhD<sup>2</sup>, Susan Pressler, PhD, RN, FAAN, FAHA<sup>3</sup>  
<sup>1</sup>University of Notre Dame, <sup>2</sup>LSUHSC School of Public Health, <sup>3</sup>Indiana University School of Nursing

## Introduction

More than 1 million cases of heart failure are diagnosed each year.<sup>1</sup> Heart failure is the single most frequent cause of hospitalization among the elderly.<sup>2</sup> Annual costs exceed \$24,000 per patient and may total \$70 billion by 2030.<sup>1,3</sup>

Cognitive decline is an important comorbidity of heart failure.<sup>4</sup> Symptoms of cognitive decline can decrease patient compliance with treatment plans and increase the risk of hospitalization. Addressing cognitive decline in the management of heart failure is a component of current clinical practice guidelines.<sup>5,6</sup>

A pilot study randomized patients with cognitive decline due to heart failure into three groups; the training program BrainHQ, puzzles, and usual care.<sup>7</sup> Costs were lower for patients in the BrainHQ group, but the difference was not statistically significant. A larger scale trial included three ways of measuring costs; electronic medical records (EMR), the Medical Resource Utilization Questionnaire (MRUQ) and the Living with Heart Failure Questionnaire (LHFQ).<sup>8</sup>



**Figure 1. Cognitive Interventions.** Patients with a MoCA level < 19 show signs of moderate to severe cognitive impairment and were thus removed from the study.


## Research Question

Do alternative cost measures yield similar results? Additionally, does the choice of cost measure affect the conclusions drawn about the cost of treatment for patients with cognitive decline due to heart failure?


## Methods

This study used disaggregated patient data from the recently completed larger trial.<sup>9</sup> Costs of medical services were compared across the cost measures and treatment groups. Stata SE 17 was used to perform Poisson regression analysis on the difference in counts.


## Hospitalizations




## Emergency Room Visits



## Clinician Visits



## Key Results



## Conclusion

Alternative cost measures do not yield similar results in the counts of use of medical services by patients with cognitive decline due to heart failure. Electronic medical records report use of more services than some patient reported questionnaires. The choice of cost measure does affect the conclusions drawn about the cost-effectiveness of treatment for patients with cognitive decline due to heart failure. Only analysis using the Medical Resource Utilization Questionnaire suggests significant differences in costs among treatment groups.

## References

1. Tsao CW, et al. Heart Disease and Stroke Statistics—2022 Update: A Report from the American Heart Association. *Circulation*. 2022;145:e153–e639.
2. Krumholz HM, et al. Mortality, hospitalizations, and expenditures for the Medicare population aged 65 years or older, 1999–2013. *JAMA*. 2015;314:355–365.
3. Urbich M, et al. A Systematic Review of Medical Costs Associated with Heart Failure in the USA (2014–2020). *Pharmacoeconomics*. 2020;38:1219–1236.
4. Becher, PM, et al. An Update on Global Epidemiology in Heart Failure. *European Heart Journal*. 2022;ehac248.
5. Liori, S, et al. Cognitive Impairment in Heart failure: Clinical implications, Tools of assessment, and Therapeutic considerations. *Heart Failure Reviews*. 2022;27:993–999.
6. Heidenreich PA, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022;145:e895–e1032.
7. Pressler, SJ, et al. Healthcare Resource Use Among Heart Failure Patients in a Randomized Pilot Study of a Cognitive Training Intervention. *Heart & Lung*. 2013;42:332–338.
8. Pressler, SJ, et al. Design and Rationale of the Cognitive Intervention to Improve Memory in Heart Failure Patients Study. *Journal of Cardiovascular Nursing*. 2018;33:344–355.
9. Pressler, SJ, et al. Randomized Controlled Trial of a Cognitive Intervention to Improve Memory in Heart Failure. *Journal of Cardiac Failure*. 2022;28:519–530.