

Local Control of Pyoderma Gangrenosum

Cameron J. Fontenot, BS ¹; Mark A. Maier, BS ¹; Nicholas Taylor, MD ²; Alison A. Smith, MD, PhD ^{2,4}; Frank Lau, MD ^{2,3}



¹ School of Medicine, Louisiana State University Health Sciences Center, New Orleans, LA ²Department of Surgery, ³Division of Plastic and Reconstructive Surgery, ⁴Division of Trauma/Critical Care Surgery cfon18@lsuhsc.edu; mmaie1@lsuhsc.edu; ntay16@lsuhsc.edu; asmi60@lsuhsc.edu; flau@lsuhsc.edu

Introduction

- Pyoderma gangrenosum (PG) is a rare, non-infectious, ulcerative dermatosis that worsens with debridement (pathergy)
- There is no consensus regarding the diagnosis, treatment, or management of PG
- Application of dehydrated amnion/chorion membrane (dHACM) following debridement has previously demonstrated successful wound healing

Objective: Characterize and compare PG wounds pre- and post-treatment with dHACM by identifying select transcriptomes

Methods

- 1. Excisional debridement of the wound and application of dHACM
- 2. Application of a split-thickness skin graft (STSG) for wound coverage within 2 weeks of treatment
- 3. RNA isolation and reverse transcriptase polymerase chain reaction of wound samples from pre- and post- treatment

Results

Course of Treatment/Outcomes

B



Figure 1. A) PG wound post-debridement B) 8 days post-operation treating PG with pathergy using second dHACM application C) 15 days post-initial operation showing wound granulation and skin graft D) 20 days post-initial operation at follow-up.

- 43-year-old man presented with a single PG wound
- The wound was on the dorsum of his left foot to the level of the fascia with a TBSA of 3.24cm²
- A subsequent round of dHACM was necessary 7 days later (Fig 1B)
- A small wound appeared lateral to the primary lesion with undermining and purpura, which is indicative of active pathergy (Fig 1B)
- A STSG was placed 15 days post-initial treatment (Fig 1C)

Summary

- PG wounds were successfully treated using dHACM and splitthickness skin grafting
- This case is part of a clinical trial for pyoderma gangrenosum (NCT05120726)

Future Directions/Conclusions

- Provide insight into the etiology, pathogenesis, and treatment of PG with dHACM
- Transcriptome analysis is on-going with additional patient enrollment

References

- 1. Snyder RJ, Ead J, Glick B, Cuffy C. Dehydrated Human Amnion/Chorion Membrane as Adjunctive Therapy in the Multidisciplinary Treatment of Pyoderma Gangrenosum: A Case Report. *Ostomy Wound Manage*. 2015;61(9):40-49.
- 2. Maverakis E, Marzano A V., Le ST, et al. Pyoderma gangrenosum. *Nat Rev Dis Prim.* 2020; 6(1): 1-19.
- 3. Koob TJ, Lim JJ, Massee M, Zabek N, Denozière G. Properties of dehydrated human amnion/chorion composite grafts: Implications for wound repair and soft tissue regeneration. *J Biomed Mater Res Part B Appl Biomater.* 2014; 102(6): 1353-1362.

Acknowledgments

- A special thank you to Elizabeth Martin, PhD for contributing to this project
- Funding is provided by the Department of Surgery at University Medical Center, New Orleans, LA
- This research project was also supported by LSU Health Sciences Center, School of Medicine